



Melanoma: Diagnosis and staging (leaflet 4 of 7)

Patient information from the British Association of Dermatologists

Produced for National Cancer Patient Information Pathways, National Cancer Action Team

Melanoma leaflets in this series, produced by the British Association of Dermatologists:

1. Prevention and risk factors
 2. Symptom awareness and early detection
 3. Referral, tests and investigations
 4. **Diagnosis and staging**
 5. Treatments
 6. Follow up care and cancer in remission
 7. Advanced disease and recurrence
-

i) Diagnosis:

Your GP will not always be able to diagnose a skin cancer, but if he or she thinks it is a possibility, they will refer you to a specialist. This specialist is usually a dermatologist or a surgeon with a special interest in pigmented lesions, and is generally based at a hospital.

The specialist will examine the area, perhaps with a special instrument (a dermatoscope), and decide whether it needs to be removed.

The only way in which the diagnosis of a melanoma can be made firmly is by removing all or part of the suspected area and looking at it under a microscope in the laboratory.

If the mole needs to be examined, the whole of the suspicious area will then be removed under a local anaesthetic (called an 'excision biopsy') and sent to the laboratory to be examined.

If the area is too large to remove easily, a sample of it (a 'biopsy') will be taken. If a melanoma is found, the biopsy specimen will provide valuable information about its type and depth that will help to plan the next step in treatment.

ii) Staging:

'Staging' is the process of determining the size of the melanoma and if / how far it has spread.

When the tumour is removed, the depth of the tumour will be examined. This is the depth from the surface of the skin at which the cancer cells are present. This is measured according to the Breslow thickness, which states to what depth in millimeters (mm) tumour cells are present at the time of surgical excision. Most melanomas are less than 1mm thick and this group are less likely to come back or require further treatment.

TNM System:

The tumour-node-metastasis system (TNM System), created by the American Joint Committee on Cancer (AJCC), is widely used system for staging of skin cancer, and expands on the Breslow scale mentioned above.

- T: Primary Tumour

The 'T' represents the thickness of the tumour in mm and whether it is 'ulcerated' (the skin is broken on the surface of the tumour).

In terms of thickness:

Tis is 'Tumour in situ': the cells are only in the epidermis, the very outermost layer of skin

T1 is less than 1.0mm

T2 is 1.01-2.0mm

T3 is 2.01-4.0mm

T4 is more than 4.0mm

This is then further defined according to whether or not the tumour is ulcerated. If it is not, it is classed as 'a' whereas if there is ulceration, it is classed as 'b'. Tumours that are ulcerated are more likely to spread. So for example, a tumour classed as T2a would be 1.01 to 2mm thick and without ulceration.

- N: Regional Lymph Nodes

The 'N' refers to whether or not the tumour has spread to the nearby lymph nodes.

N0 means there is no evidence of the melanoma cancer cells in the lymph nodes.

N1 means there are cancer cells present in one nearby lymph node.

N1a means the cancer can only be detected as cells using a microscope (microscopic metastasis).

N1b means the cancer can be felt or seen by the doctor examining you, as the lymph nodes are enlarged (macroscopic metastasis).

N2 means there are cancer cells in 2 or 3 nearby lymph nodes, or there are cancer cells in areas of skin near the tumour but not in the lymph nodes.

N2a means the cancer can only be detected as cells using a microscope (microscopic metastasis).

N2b means the cancer can be felt or seen by the doctor examining you, as the lymph nodes are enlarged (macroscopic metastasis).

N2c means the melanoma cells are present in very small areas of skin close to the tumour (satellite tumours) or has spread to skin lymphatic channels around the tumour, but without reaching the lymph nodes.

N3 means that either:

There are cancer cells in 4 or more nearby lymph nodes, or

There are cancer cells in lymph nodes that have become 'matted' together, or

The melanoma cells are present in very small areas of skin close to the tumour (satellite tumours) or the cancer has spread to skin lymphatic channels around the tumour, and also in the lymph nodes.

- M: Distant Metastasis

The 'M' signifies whether or not the cancer has metastasized (spread) to other parts of the body.

M0 means there is no evidence of the cancer having spread to distant tissues or organs

M1a means the cancer has spread to skin elsewhere on the body or to lymph nodes that are not close to the tumour

M1b means the cancer has spread to the lung

M1c means the cancer has spread to other parts of the body.

- Stage Grouping:

Once the T, N, and M are determined, they are combined to provide an overall "Stage" of the cancer – numbered 0 to 4 (often represented the Roman numerals 0, I, II, III, IV). These can be further divided as a, b or c.

Stage	Features	T	N	M
0	The cells are only present in the top layer of skin, called the 'epidermis', and have not spread to the 'dermis', the layer of skin below the epidermis. Stage 0 melanoma is also called melanoma or tumour 'in situ'	Tis	0	0

	(Tis).			
1a	Less than 1mm in the dermis. No ulceration. No spread to lymph nodes or other parts of the body.	T1a	0	0
1b	Less than 1mm in the dermis and ulcerated or 1mm to 2mm and not ulcerated. No spread to lymph nodes or other parts of the body.	T1b T2a	0 0	0 0
2a	1 - 2 mm and ulcerated or 2 - 4mm and not ulcerated. No spread to lymph nodes or other parts of the body.	T2b T3a	0 0	0 0
2b	2 - 4mm and ulcerated or More than 4mm and not ulcerated. No spread to lymph nodes or other parts of the body.	T3b T4a	0 0	0 0
2c	More than 4mm and ulcerated. No spread to lymph nodes or other parts of the body.	T4b	0	0
3a	1 - 4mm and not ulcerated. Cancer cells present in one, two or three nearby lymph nodes. The cancer in the lymph nodes can only be detected as cells using a microscope, not felt or seen by the doctor examining you (the lymph nodes are not enlarged.) No spread to other parts of the body.	T1-4a T1-4a	1a 2a	0 0
3b	1 – 4mm and ulcerated. Cancer cells present in one, two or three nearby lymph nodes. The cancer in the lymph nodes can only be detected as cells using a microscope, not felt or seen by the doctor examining you (the lymph nodes are not enlarged.) No spread to other parts of the body.	T1-4b T1-4b	1a 2a	0 0

	<p>or</p> <p>1 - 4mm and not ulcerated.</p> <p>Cancer cells present in one, two or three nearby lymph nodes. The cancer in the lymph nodes can be felt or seen by the doctor examining you (the lymph nodes are enlarged.)</p> <p>or</p> <p>1 – 4mm, either ulcerated or not ulcerated. Cancer cells present in very small areas of skin close to the tumour (satellite tumours) or has spread to skin lymphatic channels around the tumour, but without reaching the lymph nodes.</p> <p>No spread to other parts of the body.</p>	<p>T1-4a</p> <p>T1-4a</p> <p>T1-4a/b</p>	<p>1b</p> <p>2b</p> <p>2c</p>	<p>0</p> <p>0</p> <p>0</p>
3c	<p>1 - 4 mm and ulcerated. Cancer cells present in 1 to 3 nearby lymph nodes. The cancer in the lymph nodes can be felt or seen by the doctor examining you (the lymph nodes are enlarged.)</p> <p>1 – 4mm, either ulcerated or not ulcerated. Cancer cells present in 4 or more nearby lymph nodes, or there are cancer cells in lymph nodes that have become ‘matted’ together, or the melanoma cells are present in very small areas of skin close to the tumour (satellite tumours) or the cancer has spread to skin lymphatic channels around the tumour and is also present in the nearby lymph nodes.</p> <p>No spread to other parts of the body.</p>	<p>T1-4b</p> <p>T1-4b</p> <p>Any T</p>	<p>1b</p> <p>2b</p> <p>N3</p>	<p>0</p> <p>0</p> <p>0</p>
4	<p>The melanoma cells have spread elsewhere in the body, away from the primary tumour and nearby lymph nodes, such as to organs like the brain or lung, or to distant lymph nodes.</p>	<p>Any T</p>	<p>Any N</p>	<p>M1 (a,b or c)</p>

For more information, please contact:

British Association of Dermatologists

4 Fitzroy Square, London W1T 5HQ

website: www.bad.org.uk

tel: 0207 383 0266

© British Association of Dermatologists 2009