



PHOTODYNAMIC THERAPY

What are the aims of this leaflet?

This leaflet has been written to help you understand more about photodynamic therapy - what it is, what is involved, what the potential side effects are, and where to find out more about it.

What is photodynamic therapy?

Photodynamic therapy (PDT) is a technique for treating skin cancers and sun-damaged skin which might one day turn cancerous (pre-cancers). In PDT a light beam activates a cream which has been applied to the lesion (affected area of skin). This treatment kills the abnormal cells in the skin.

How does photodynamic therapy work?

PDT involves the use of a light-sensitive chemical (called a *photosensitiser*). This photosensitiser is, by itself, harmless. When light of a certain wavelength (usually red light) is shone onto skin to which the photosensitiser has been applied, the photosensitiser is activated. This causes changes in the oxygen molecules within the sun-damaged skin cells. These ~~%excited+~~ excited oxygen molecules kill the cells. Only the area of skin exposed to the light source will be affected and inflamed; after the inflammation clears out it will be cured.

What skin conditions can be treated with photodynamic therapy?

PDT can be used to treat various skin conditions including:

- Thin [basal cell carcinomas](#)
- [Bowen's disease](#) (*in situ* squamous cell carcinoma), a pre-cancer.
- [Actinic \(solar\) keratoses](#) - early sun-damage, a pre-cancer

What does photodynamic therapy for skin lesions involve?

PDT is an outpatient procedure, and will be done by your doctor or nurse. It takes several hours to complete. The first step is to apply a cream containing the photosensitiser to the area that needs to be treated. If necessary, any loose scale or crusts will be removed first. A dressing will then be applied over the cream and you will be asked to return in about 3 hours. This wait is to allow the photosensitiser to be absorbed and to be converted into the active chemical by the skin. The cream is then wiped off and the area cleaned. Then, a bright coloured light is shone onto the treatment area for approximately 10 to 45 minutes (the precise time will be determined by your doctor or nurse). After the treatment has been completed, a dressing will be applied for a minimum of 2 days, to prevent any further exposure to light.

How should the treated area be cared for?

Your doctor will explain how you should care for the treated areas. It is usually suggested that, after the dressing has been removed, you can wash, bathe or shower as usual. Do not rub the treated area, but gently pat it dry. Within a few days, a scab will form, and healing will take several weeks (depending on which part of the body has been treated). Care must be taken not to scratch the area or accidentally dislodge the scab during the healing process. The use of a suitable sunscreen (SPF 30) following the procedure, especially during outdoor activities, is important.

What reasons might prevent you having photodynamic therapy?

- PDT is not recommended if you are pregnant.
- The PDT cream may contain peanut oil, so tell your doctor and nurse if you are allergic to peanuts.
- PDT is not recommended if you have porphyria (a light-sensitive disorder, where the patient is producing the same photosensitiser used in PDT and is present in the blood, skin and other tissues of the body).

Do I need to avoid anything whilst having photodynamic therapy?

Care should be taken not to get the dressing wet when bathing or showering. It is also best to avoid swimming until the treated area is fully healed.

What are the potential side effects of photodynamic therapy?

The short-term side effects of phototherapy include:

- *Pain*. When the red light is shone onto the skin, the treated area may hurt. If it is too uncomfortable, your doctor or nurse may suggest pausing treatment for a while, or a local anaesthetic injection may be recommended. After completion of treatment, discomfort and itching may last for a few days, and may require pain-killers.
- *Inflammation*. The treated area may initially become pink and puffy, and may ooze a little: this is a normal reaction. It settles within a few days.
- *Blistering and ulceration*. The treated area may occasionally blister or ulcerate.
- *Infection*. If the treated area becomes red, swollen and painful, an infection may have developed, and you should contact your doctor.

Potential long-term side effects of phototherapy include:

- *Scarring*. There may be some scarring after PDT.
- *Colour change*. The skin may be left darker or paler after PDT.
- Treatment may not be effective, or the condition may come back again. If this happens, you may be offered further PDT, or an alternative type of treatment may be recommended.

Top sun safety tips

- Protect the skin with clothing, including a hat, T shirt and UV protective sunglasses
- Seek shade between 11 am and 3 pm when it's sunny
- Use a sunscreen of at least SPF 30 which also has high UVA protection
- Keep babies and young children out of direct sunlight
- The British Association of Dermatologists recommends that you tell your doctor about any changes to a mole - if your GP is concerned about your skin, make sure you see a Consultant Dermatologist (on the GMC register of specialists). Your GP can refer you via the NHS.

Where can I get more information about photodynamic therapy?

You can find more information about photodynamic therapy on the following websites:

www.cancerbackup.org.uk/Treatments/Othertreatments/Photodynamictherapy
www.cancerhelp.org.uk
<http://dermnetnz.org/procedures/photodynamic-therapy.html>

This leaflet aims to provide accurate information about the subject and is a consensus of the views held by representatives of the British Association of Dermatologists: its contents, however, may occasionally differ from the advice given to you by your doctor.

This leaflet has been assessed for readability by the British Association of Dermatologists' Patient Information Lay Review Panel

**BRITISH ASSOCIATION OF DERMATOLOGISTS
PATIENT INFORMATION LEAFLET
PRODUCED SEPTEMBER 2009
REVIEW DATE SEPTEMBER 2012**